

Binaries among Ap and Am stars

P. North¹, N. Ginestet², J.-M. Carquillat², F. Carrier³ and S. Udry³

¹ *Institut d'Astronomie de l'Université de Lausanne,
CH-1290 Chavannes-des-Bois, Switzerland*

² *Observatoire Midi Pyrénées, 14 avenue Edouard Belin, F-31400 Toulouse,
France*

³ *Observatoire de Genève, CH-1290 Sauverny, Switzerland*

Abstract. The results of long-term surveys of radial velocities of cool Ap and Am stars are presented. There are two samples, one of about 100 Ap stars and the other of 86 Am stars. Both have been observed with the CORAVEL scanner from Observatoire de Haute-Provence (CNRS), France.

The conspicuous lack of short-period binaries among cool Ap stars seems confirmed, although this may be the result of an observational bias; one system has a period as short as 1.6 days. A dozen new orbits could be determined, including that of one SB2 system. Considering the mass functions of 68 binaries from the literature and from our work, we conclude that the distribution of the mass ratios is the same for the Bp-Ap stars than for normal G dwarfs.

Among the Am stars, we found 52 binaries, i.e. 60%; an orbit could be computed for 29 of them. Among these 29, there are 7 SB2 systems, one triple and one quadruple system. The 21 stars with an apparently constant radial velocity may show up later as long-period binaries with a high eccentricity. The mass functions of the SB1 systems are compatible with cool main-sequence companions, also suggested by ongoing spectral observations.

Key words: Stars: binaries: spectroscopic – Stars: chemically peculiar – Stars: evolution

1. Introduction

The question here is that of the possible influence of a binary companion upon chemical peculiarity: does an Ap or Am star “need” a companion, or conversely, does it need to be single? If there is a companion, what is its nature? Finally, fundamental parameters like mass, radius and age are better constrained for the components of binary stars than for isolated ones.

The first systematic search for spectroscopic binaries among Bp-Ap stars was made by Abt & Snowden (1973) on a sample of 62 northern bright stars. They found a rate of binaries of only 20 percent among the magnetic Ap stars and of 43 percent among HgMn stars (the rate among normal stars is 47 ± 5 percent among normal B0 to M stars according to Jaschek & Gomez 1970). Aikman

(1976) studied 80 HgMn stars (instead of 15 for Abt & Snowden) and confirmed a normal rate with 49 percent of binaries. The most recent and complete review is that of Gerbaldi *et al.* (1985), while North (1994) published some new results on cool Ap stars obtained with the CORAVEL spectrovelocimeter.

On the basis of a sample of 25 Am stars, Abt (1961) concluded that all are binaries, while Abt & Levy (1985) confirmed the high rate of binaries in a larger sample but also found some stars to be apparently single. For both Ap and Am stars, only bright objects had been examined, so it was judged worthwhile to extend the survey to fainter and more numerous ones.

This contribution is kind of a progress report on a survey of cool Ap and Am stars made with the CORAVEL scanner in the northern hemisphere.

2. Samples and observations

All observations have been made at Observatoire de Haute-Provence (CNRS), France, with the CORAVEL spectrovelocimeter (Baranne *et al.* 1979) attached to the 1m Swiss telescope. The CORAVEL scanner is optimized for cool stars, since the mask yielding the correlation function was made from the spectrum of the yellow giant Arcturus. Nevertheless, even stars with as hot a type as A0 may provide an autocorrelation dip, provided they rotate slowly enough and have enhanced metallic lines, which is precisely the case of many (though not all) Ap and Am stars.

2.1. Cool (SrCrEu) Ap stars

Some well-known magnetic Ap stars had been already measured by Dr. Michel Mayor and coworkers as early as 1980, but a more complete sample has been defined in 1985 from the following criteria:

- Ap classification from e.g. the compilation of Bertaud & Floquet (1974)
- Ap stars having Geneva colours
- $T_{\text{eff}} \lesssim 10000$ K according to Geneva photometry
- declination $\delta \geq -20^\circ$
- $V \leq 8.6$ (some fainter stars included as well)

This makes a total of 160 stars, but 48 of them proved to yield no autocorrelation dip because of fast rotation. Indeed, the dip becomes too flat when $v \sin i \gtrsim 30 \text{ km s}^{-1}$, so that our sample is severely biased against high $v \sin i$. This implies that the sample is also biased against periods shorter than about 4 days, if synchronism is assumed and if $i \sim 90^\circ$ (assuming also the orbital and equatorial planes to be coplanar); systems with shorter periods may of course be observed, provided the inclination i is small enough. Among the 112 stars left, 13 are more probably Am than Ap stars because some early, unreliable classification sources were used, and a few more had a so wide and shallow correlation dip that they were observed only once or twice. Therefore, we are left with 95 objects measured ≥ 3 times (typically 6-8 times) over a time span of 12 to 17 years.

2.2. Am stars

Systematic observations aimed at studying the binaries began in 1992. The sample was defined as including

- northern objects
- metallic type F2 or later according to the compilation by Hauck (1986); some objects from Bidelman (1988) and from Hynek (1938) were also included
- spectroscopic binaries with known orbits were excluded from the sample

As for the sample of Ap stars, this one is biased against high $v \sin i$ (therefore against very short orbital periods). But it is also biased, to some extent, against large V_r amplitude, short period spectroscopic binaries, because objects with known and published orbits – which are more easily discovered when the period is short – are excluded. There are 230 objects in this sample, among which 158 have been observed to date. However, as many as 72 stars did not yield any autocorrelation dip, probably because of their high rotational velocity, so we are left with 86 objects.

3. Results

3.1. Ap stars

Because of the inhomogeneous distribution of the elemental abundances on the surface of these stars, many of them are spectrum variables and present radial velocity variations due to rotation and abundance patches. This problem was avoided by Abt & Snowden (1973) by using hydrogen rather than metallic lines, but this is of course not possible with CORAVEL. We are forced to identify these stars *a posteriori*, which is possible at least in part, thanks to some typical characteristics like wide and shallow correlation dip, short timescale (a few days) and relatively low-amplitude variability ($\sim 20 \text{ km s}^{-1}$) leading to a very small “mass function” if one tries to fit an orbit. Taking this into account, one has 2 SB2 and 24 certain SB1 binaries among all 95 stars, which makes 27 percent. This percentage is smaller than that of normal stars ($47 \pm 5 \%$).

What is more interesting is that we found several long-period binaries but relatively few short period ones, which seems to confirm the lack of tight orbits among Ap stars in spite of the bias we have against periods shorter than 3-4 days. This is entirely consistent with the results of Gerbaldi et al. (1985), who found no orbital period shorter than about 3 days. However, we have found HD 200405 to be a spectroscopic binary with $P_{\text{orb}} = 1.635$ days, which is the shortest orbital period known among Ap stars. Figure 1a shows the e vs. $\log P$ diagram for all known spectroscopic binaries with an Ap stars, including 9 new orbits determined with CORAVEL. This diagram resembles very closely that of normal stars (see Fig. 1 of Gerbaldi et al. 1985), the only difference being the cutoff at about 0.5. This cutoff suffices to explain the peculiar distribution of eccentricities of the Ap binaries shown in Fig. 7 of Gerbaldi et al. (1985): the lack of low

eccentricities is linked with the lack of very short periods among Si and SrCrEu binaries; HgMn stars tend to have shorter periods than the latter, which also explains why there are more circular orbits among them. Thus, the distribution of eccentricities does not appear to be peculiar in itself but only reflects the distribution of orbital periods. Figure 1b shows the cumulative distribution of




Figure 1. **a:** e - $\log P$ diagram for Ap stars. Key to symbols: open triangles=HgMn, black triangles=He weak, open squares=SrCrEu, black squares=Si. **b:** Cumulative distribution of all Ap binaries (including HgMn) compared with a simulated one.

the mass functions of all binaries hosting an Ap star (including the HgMn, but the result would be the same without them). The continuous line is the result of a Monte-Carlo simulation where the mass ratio is a gaussian, as proposed by Duquennoy & Mayor (1991) for the nearby G dwarfs; the mass function of the primaries is Salpeter’s one multiplied by the relative mass function of Ap stars given by North (1994). The observed and simulated distributions are in perfect agreement, suggesting that the unseen companion of binary Ap stars has nothing exotic; in particular, it is generally not a white dwarf.

3.2. Am stars

We have found as yet 52 binaries in our sample of 86 objects, i.e. 60 percent. 29 orbits could be determined for 19 SB1 systems, 7 SB2 ones, one triple and one quadruple system. 13 stars have not been measured often enough to conclude about their possible V_r variability, but the remaining 21 seem to be constant, hence single; however, additional observations may show them to be variable, if they belong to long-period, highly eccentric systems. Figure 2 shows the e - $\log P$ diagram of our 29 orbits, which closely resembles that of the sample of Abt & Levy (1985). Here too, there is a lack of very short periods ($P_{\text{orb}} \lesssim 1.0 - 1.5$ days): this can be understood from the fact that no Am star has an equatorial velocity larger than 100 km s^{-1} , while synchronism in such systems would force the primary to rotate faster. The distribution of periods is very similar to that of an earlier sample (Ginestet et al. 1982), with a maximum between 5 and 10 days. But, since observational biases tend to favour such short period systems, it remains necessary to accumulate observations on a long timescale. On Figure 2

are also shown the V_r amplitudes K_1 vs. $v \sin i$ of the primaries for all 29 systems with a known orbit, as well as the $v \sin i$ of the 21 stars with an apparently constant V_r . The upper envelope of the points is traced by SB2 systems with $M_2/M_1 \sim 1$, 4 of which at least are already synchronized. The mass functions of SB1 Am binaries are compatible with a main sequence cooler secondary, as is the case of Ap binaries.




Figure 2. *left:* e - $\log P$ diagram for Am stars. Key to symbols: asterisks=SB1 systems, full dots=SB2 systems. *right:* K_1 vs. $v \sin i$; same symbols as above; the open dots are for stars with a constant V_r . The mass ratio is given for the SB2 systems.

References

- Abt, H.A.: 1961, *Astrophys. J., Suppl. Ser.* **6**, 37
 Abt, H.A., Levy, S.G.: 1985, *Astrophys. J., Suppl. Ser.* **59**, 229
 Abt, H.A., Snowden, M.S.: 1973, *Astrophys. J., Suppl. Ser.* **25**, 137
 Aikman, G.C.L.: 1976, *Publ. Dominion Astrophys. Obs.* **14**, 379
 Baranne, A., Mayor, M., Poncet, J.-L.: 1979, *Vistas Astron.* **23**, 279
 Bertaud, C., Floquet, M.: 1974, *Astron. Astrophys., Suppl. Ser.* **16**, 71
 Bidelman, W.P.: 1988, *Publ. Astron. Soc. Pac.* **100**, 1084
 Duquennoy, A., Mayor, M.: 1991, *Astron. Astrophys.* **248**, 485
 Gerbaldi, M., Floquet, M., Hauck, B.: 1985, *Astron. Astrophys.* **146**, 341
 Ginestet, N., Jaschek, M., Carquillat, J.M., Pedoussaut, A.: 1982, *Astron. Astrophys.* **107**, 215
 Hauck, B.: 1986, *Astron. Astrophys., Suppl. Ser.* **64**, 21
 Hynek, J.A.: 1938, *Contrib. Perkins Obs.* No 10
 Jaschek, C., Gomez, A.E.: 1970, *Publ. Astron. Soc. Pac.* **82**, 809
 North, P.: 1994, in *25th meeting and workshop of the European Working Group on CP stars*, eds: I. Jankovics and I.J. Vincze, Gothard Astrophysical Observatory, Szombathely, Hungary, 3
 Wade, G.A., North, P., Mathys, G., Hubrig, S.: 1996, *Astron. Astrophys.* **314**, 491