

Determination of T_{eff} and $\log g$ for A0 type stars and its impact on the interpretation of the HR diagram with the HIPPARCOS results

M. Gerbaldi^{1,2}, R. Faraggiana³, R. Burnage⁴, F. Delmas¹,
A. Gomez⁵ and S. Grenier⁵

¹ Institut d'Astrophysique - 98bis, Bd. Arago - 75014 Paris - France

² Université de Paris-Sud XI

³ Dipartimento di Astronomia - v. Tiepolo 11 - 34131 Trieste - Italy

⁴ Observatoire de Haute-Provence - 04870 Saint Michel l'Observatoire - France

⁵ DASGAL, Observatoire de Meudon - 92150 Meudon - France

Abstract. The determination of the “Fundamental Parameters” T_{eff} and $\log g$ for a set of dwarf A0-type stars is discussed in terms of consistency when comparing these values determined through different methods. The position of these stars in the HR diagram are discussed, taking into account the HIPPARCOS data. A large number of binary stars with components of similar spectral types has been found from this spectroscopic survey.

Key words: Stars: fundamental parameters – Stars: A0-type – HIPPARCOS – HR diagram

1. Introduction

Our purpose is to analyse a set of A0-type, non-giant stars, in order to define a sample to be used as *standard stars* for further studies. These stars have parallaxes measured from the HIPPARCOS experiment and their position in the HR diagram will be used to derive their evolutionary status. The spectra obtained in the H_{γ} region are used to detect peculiarities as well as to check the fit with theoretical spectra computed with T_{eff} and $\log g$ values determined from calibrations of colour indices.

2. Selection of the sample

All stars with a spectral type A0V in the Bright Stars Catalogue (1982 ed.) (hereafter BSC), excluding those recognized as peculiar, shell stars and SB2 have been selected; this sample contains 230 objects. Spectroscopic data refer

to a second sample of A0 stars brighter than the 7th magnitude extracted from a larger set of A-type stars defined in the framework of our ESO Key programme (Gerbaldi et al. 1989). Slightly more than 70 A0 type stars were extracted from this set, for which spectra were obtained in the spectral range 4200 - 4500 Å using the Echelec spectrograph attached to the 1.52-m ESO telescope. All the observations were done at the European Southern Observatory (ESO), La Silla, Chile. The already known peculiar stars, Ap and Am, were *a priori* excluded from the sample.

3. Atmospheric Parameters

The atmospheric parameters T_{eff} and $\log g$ were derived from calibrated colour indices because we had not enough data to derive T_{eff} from the Infrared flux method (Blackwell et al. 1980). Very few attention has been given up to now to the dwarf stars in the range of spectral types A0 – A3 where the Balmer line profiles vary with both T_{eff} and $\log g$.

Any calibration of colour indices in terms of T_{eff} and $\log g$ rely upon :

- synthetic indices

These are affected by the ingredients used to compute the atmosphere models and then the synthetic spectra (chemical abundances, convection...)

The synthetic indices are used to define the *zero point* by comparing their values to those of some standard stars.

- *well determined* values of T_{eff} and $\log g$

The most commonly used T_{eff} values are those deduced by Code et al. (1976) using the angular diameters measured by Hanbury-Brown et al. (1974) and the energy distribution from the UV to the Infrared. The $\log g$ values are obtained from binary systems.

Very recently new measurements of the stellar angular diameters have been done in Australia (see Booth, 1997), but they have not been incorporated into the re-calibration of any photometric system. The accuracy of the Infrared flux method to determine T_{eff} has been extensively discussed by Mégessier (1994, 1995, 1997a, 1997b).

The best sources of information for T_{eff} and $\log g$ for our sample of early type stars, are the uvby β and Geneva photometric systems. Several calibrations of the uvby β system exist; we shall use that by Moon & Dworetsky (1985) (hereafter quoted MD) which has been tested and refined since more than a decade.

The MD calibration relies upon synthetic colours and indices from Relyea and Kurucz (1978), Philip and Relyea (1979), Schmidt (1979) and Kurucz (1979). The zero point has been adjusted by means of stars with *well determined* T_{eff} and $\log g$ values. The photometric calibration for B-type stars has been re-evaluated by Castelli (1991). Tests on this calibration done by Balona (1994) and by Napiwotzki et al. (1993) have shown that no important modification of the MD calibration is needed for early A-type stars. Napiwotzki et

al. (1993) suggested a small correction to be done on $\log g$ for $T_{\text{eff}} \geq 9000$ K. Smalley and Dworetsky (1995) re-evaluated the *fundamental* values of T_{eff} as determined by Code et al. (1976) using recent flux measurements; no significant differences were found by these authors. To conclude, up to now, no revision of the MD calibration of the uvby β is required for dwarf A0 stars.

The calibration of the Geneva photometry for early-type stars is very recent (Künzli et al. 1997); this system is quite independent from the previous one in the sense that it does not include any narrow band centered on a Balmer line; moreover, its calibration is done using recent Kurucz flux computations and it is based on a different and larger set of standard stars.

It is therefore interesting to compare the results obtained from these two systems and to analyse the coherence of the parameters obtained from different aspects of the emitted stellar flux.

4. T_{eff} and $\log g$ accuracy

4.1. Influence of the colour excess E(b-y) on T_{eff} , $\log g$

Before computing the atmospheric parameters, the observed colour indices must be de-reddened. For the bright stars of our sample, their visual magnitude being less than 7.0 mag, the reddening is expected to be null or negligible.




Figure 1. For the stars with $2.869 \leq \beta \leq 2.880$, relation between the T_{eff} computed with the “early group” choice and the T_{eff} computed with the “intermediate” choice

Even though the reddening is expected to be negligible, we tested it; we used the programme UVBYBETA by Moon (1985) in order to check the consistency of the observed values in term of “normality” of the star. We know that any “blueing” can be interpreted as due to a Bp-Ap effect, and any strong reddening requires to have a close look at the spectrum. The determination of the amount of reddening is based on an empirical calibration of the uvby β system, so it is free from any atmospheric model. Jordi et al. (1997) have discussed the reddening law of Hilditch (1983) adopted by Moon (1985); they have shown that Moon overestimates the E(b-y) value compared to another de-reddening law : the one by Grosbol (1978). Due to the fact that the stars of our sample have a small value of their colour excess E(b-y), the error made by using one or the other reddening law will be small.

We have shown (Gerbaldi et al. 1997) that a moderate reddening of E(b-y)=0.01 produces a difference in T_{eff} of 200K (see Figure 1 in Ref. cited); on log g determination, the effect is negligible (see Figure 2 in Ref. cited). The stars which are outside the mean relation have values of the parameters a_0 and r which are outside the limits of the grids used by the MD programme; it is a numerical effect which produces an incorrect result.

In fact a small non-zero value for E(b-y) can arise from uncertainties on the colours as well as on the *group choice*.




Figure 2. For the stars $2.880 \leq \beta \leq 2.890$, relation between the T_{eff} computed with the “intermediate” choice and the T_{eff} computed with the “early group” choice

4.2. Effect of the group choice on T_{eff} and $\log g$

Dereddening procedures for A0-type stars require to assign these stars to one of these groups:

- early group : $2.59 \leq \beta \leq 2.88$
- intermediate group $2.87 \leq \beta \leq 2.93$

In Moon's (1985) programme UVBYBETA the information concerning the spectral type is added. The assignment to a group is sometimes ambiguous, errors in group selection cannot be avoided and this may produce uncertain dereddened colours. We have estimated the effect on T_{eff} , $\log g$ of a wrong group selection, which can be due to an observational error on the value of β . To select the stars to do such a test, we have considered an error on β of ± 0.01 , so the β limits are 2.869 and 2.890.




Figure 3. For the stars with $2.869 \leq \beta \leq 2.880$, and $2.880 \leq \beta \leq 2.890$, relation between the color excess $E(b-y)$ computed in the normal group of the star according to the value of β and $E(b-y)$ computed in the other group. The number written next to some points indicates the number of stars having the same values

From our sample of AOV stars we have 28 stars with $2.869 \leq \beta \leq 2.880$ and 22 stars in the range $2.880 \leq \beta \leq 2.890$. For each selection of stars we computed T_{eff} , $\log g$ twice, that is in each group: early and intermediate. The results are displayed on Figs. 1 and 2. For the stars which normally belong to the *early* group the computations in the *intermediate* group give a difference ranging between -70 K and 80 K. For the stars which normally belong to the *intermediate* group the computations in the *early* group give a difference ranging

between -160 K and 60 K. The effect on the colour excess $E(b-y)$ is at most of ± 0.02 (Fig. 3).

5. Comparison between T_{eff} and $\log g$ determined from uvby β and Geneva Photometry

Since there is no programme to compute independently the colour excess in the Geneva photometric system, we have compared the T_{eff} and $\log g$ values only for stars with no reddening. We consider, as above, normal A0V stars from the BSC, but with a further restriction such as : $-0.015 \leq E(b-y) \leq 0.015$. There are slightly more than 140 objects. For both photometric calibrations the parameters were computed using $E(b-y)=0$.

We note (see Fig. 4) that the values of T_{eff} computed with the MD calibration are systematically higher than those derived from the Geneva photometry calibration by Künzli et al. (1997). The mean value of the difference between these two T_{eff} is -70 K with a rms of 137 K.




Figure 4. For the stars with $-0.015 \leq E(b-y) \leq 0.015$, relation between the T_{eff} computed with the MD calibration and the difference between the T_{eff} derived from the calibration of the Geneva photometric system and the T_{eff} from MD

There is also a small systematic effect on the $\log g$ determination: for $\log g \leq 3.9$ the value from Geneva photometry is higher than that from the MD calibration of uvby β colours and it is the reverse for $\log g > 3.9$.




Figure 5. The HR diagram; the evolutionary tracks from the top corresponds respectively to $3.16 M_{\odot}$, $2.51 M_{\odot}$ and $1.99 M_{\odot}$. The various symbols correspond to the following values of the gravity : hexagon $3.3 < \log g \leq 3.8$; cross : $3.8 < \log g \leq 4.15$; triangle: $4.15 < \log g \leq 4.3$. The error bars are those given by the error on the parallax measurements. The luminosity class is written.

6. Analysis of the spectroscopic observations

We had spectra at a resolution of 28000 for slightly more than 70 stars, most of them being classified A0V and the few remaining ones having a luminosity class IV or III, or no luminosity class at all. The reduction of the spectra was done as described by Burnage & Gerbaldi (1990, 1992). Synthetic spectra were computed using the Kurucz solar-abundance models (1993), and for the atmospheric parameters as computed by the MD programme. A careful comparison of the synthetic and the observed spectra allows us to detect a small number of:

- hot Am star
 - shell star
 - stars with distorted line profiles which are obviously a signature of binarity.
- After such a selection we find that about half of the stars in the original sample can be considered normal A0 dwarf stars.

It is striking to note how many binary stars with similar components have been found from this spectroscopic survey.

7. Conclusion : the HR diagram

With the subsample of normal stars selected above, we have plotted the HR diagram (Fig. 5). In abscissa, T_{eff} is that from the MD calibration, and M_V is derived from the HIPPARCOS parallax data. We remark that there is no direct relation between the luminosity class given by the MK classification and the M_V value; stars belonging to luminosity class V cover a broad range of M_V . The evolutionary tracks by Claret & Giménez (1992) are overplotted on this diagram.

The spread on Fig. 5 points out the difficulty to calibrate the spectral type A0 and luminosity class V in terms of M_V . This spread is so large that it prevents the detection of binaries simply from an anomalous position in the HR diagram of stars for which only the spectral classification is known.

References

- Balona, L.: 1994, *Mon. Not. R. Astron. Soc.*, **268**, 119
 Blackwell, D.E., Petford, A.D., Shallis, M.J. et al.: 1980, *Astron. Astrophys.*, **82**, 249
 Booth, A.J.: 1997, *IAU Symposium* **189**, eds.: T.R. Bedding et al., Kluwer, 147
 Burnage, R., Gerbaldi, M.: 1990, *2nd ESO/ST-ECF Data Analysis Workshop*, eds.: D. Baade and P. Grosbol, ESO Garching, 137
 Burnage, R. and Gerbaldi, M.: 1992, *4nd ESO/ST-ECF Data Analysis Workshop*, eds.: P. Grosbol, R.C.E. de Ruijsscher, ESO Garching, 159
 Castelli, F.: 1991, *Astron. Astrophys.*, **251**, 106
 Claret, A., Giménez, A.: 1992, *Astron. Astrophys., Suppl. Ser.*, **96**, 255
 Code, A.D., Davis, J., Bless, R.C., Hanbury-Brown, R.: 1976, *Astrophys. J.*, **203**, 417
 Gerbaldi, M., Mayor, M.: 1989, *The Messenger*, **56**, 12
 Gerbaldi, M., et al.: 1997, *Highlight in Astronomy, 23rd GA of the IAU, JD16*, in press
 Grosbol, P.: 1978, *Astron. Astrophys., Suppl. Ser.*, **32**, 409
 Hanbury-Brown, R., Davis, J., Allen, L.R.: 1974, *Mon. Not. R. Astron. Soc.*, **167**, 121
 Hilditch, R.W., Hill, G., Barnes, J.V.: 1983, *Mon. Not. R. Astron. Soc.*, **204**, 241
 Jordi, C., Masana, E. et al.: 1997, *Astron. Astrophys., Suppl. Ser.*, **123**, 83
 Künzli, M., North, P. et al.: 1997, *Astron. Astrophys., Suppl. Ser.*, **122**, 51
 Kurucz, R.L.: 1979, *Astrophys. J., Suppl. Ser.*, **40**, 1
 Kurucz, R.L.: 1993, CD-Rom 18
 Mégezzier, C.: 1994, *Astron. Astrophys.*, **289**, 202
 Mégezzier, C.: 1995, *Astron. Astrophys.*, **296**, 771
 Mégezzier, C.: 1997a, *IAU Symposium* **189**, eds.: T.R. Bedding et al., Kluwer, 61
 Mégezzier, C.: 1997b, *IAU Symposium* **189**, eds.: T.R. Bedding et al., Kluwer, 153
 Moon, T.T.: 1985, *Comm. from the Univ. of London Obs.*, **78** and Revision in: 1985,
 private comm.
 Moon, T.T., Dworetsky, M.M.: 1985, *Mon. Not. R. Astron. Soc.*, **217**, 305
 Napiwotzki, R., Schönberner, D., Wenske, W.: 1993, *Astron. Astrophys.*, **268**, 653
 Philip, A.G.D., Relyea, L.J.: 1979, *Astron. J.*, **84**, 1743
 Relyea, L.J., Kurucz, R.L.: 1978, *Astrophys. J., Suppl. Ser.*, **37**, 45
 Schmidt, E.G.: 1979, *Astron. J.*, **84**, 1739
 Smalley, B., Dworetsky, M.M.: 1995, *Astron. Astrophys.*, **293**, 446