

The hot spot in HT Cas during its Superoutburst

K. Bąkowska^{1,2} and A. Olech¹

¹ *Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences
ul. Bartycka 18, 00-716 Warszawa, Poland*

² *Astronomical Observatory Institute, Faculty of Physics,
A. Mickiewicz University, ul. Słoneczna 36, 60-286 Poznań, Poland*

Received: October 7, 2013; Accepted: December 30, 2013

Abstract. We present the analysis of eclipses observed in the dwarf nova HT Cas during its superoutburst in November 2010. Detection of a hot spot was confirmed.

Key words: Stars: individual: HT Cas - binaries: close - novae, cataclysmic variables

1. Introduction

Cataclysmic variables are close binary stars containing a white dwarf (the primary) and a main-sequence star (the secondary). In non-magnetic systems the material flows from the secondary and forms a luminous accretion disk around the primary. The hot spot is the location where the matter hits the accretion disk (for revision Hellier, 2001).

In 2010, after 25 years of quiescence and normal outbursts HT Cas went into a superoutburst. We present an analysis of two eclipses observed during this unique phenomenon. Reconstruction of the hot spot was made with the decomposition method. Details and beautiful examples of an application of this method for cataclysmic variables UX UMa, Z Cha and OY Car were presented by Smak (1994 a, 1994 b, 2007, 2008 b).

2. Results

The superoutburst in HT Cas from November 2010 lasted 11 nights (2455504 - 2455514 HJD, below marked as 504 - 514 HJD). In Fig. 1 on the left panel there are two light curves with eclipses observed in HT Cas. On the left top panel around phases $-0.1 < \phi < -0.05$ and $0.05 < \phi < 0.1$ one can see the beginning and the end of the eclipse of the hot spot, and this fact is confirmed by the reconstruction of the hot spot presented on the top right panel. On the bottom left panel we present the eclipse where the contribution of the hot spot is not so obvious to observe and the reconstruction presented on the right bottom panel confirms that the luminosity of the hot spot is indeed noticeably smaller.

The manifestation of the hot spot was explicit during the second night of superoutburst (505 HJD). The amplitude of the hot spot luminosity was $l_{s,max} =$




Figure 1. Left panel: observational data, on top the eclipse from the 2nd night of the superoutburst (505 HJD), bottom panel: the eclipse from the 6th night of the superoutburst (509 HJD). Right panel: reconstruction of the hot spot obtained with the decomposition method.

14.7% and the phase corresponding to the amplitude was $\phi_{max} = -0.072$. Four nights later, the hot spot luminosity was clearly smaller with the amplitude of $l_{s,max} = 3.85\%$ and the phase $\phi_{max} = -0.052$.

Manifestation of the hot spot in HT Cas during its superoutburst confirms that mass transfer is strongly enhanced. This is in full agreement with the EMT (Enhanced Mass Transfer) model proposed by Smak (1991, 2004, 2008 a). The analysis made with the decomposition method of eclipses observed during the November 2010 superoutburst in HT Cas gives the newest observational evidence for the EMT model.

Acknowledgements. We want to thank Prof. Smak for his thoughtful comments about the decomposition method. Project was supported by Polish National Science Center grants awarded by decision DEC-2012/07/N/ST9/04172 for KB.

References

- Hellier, C.: 2001, *Cataclysmic Variable Stars*, Springer, Chichester UK
 Smak, J.: 1991, *Acta Astron.* **41**, 269
 Smak, J.: 1994 a, *Acta Astron.* **44**, 45
 Smak, J.: 1994 b, *Acta Astron.* **44**, 59
 Smak, J.: 2004, *Acta Astron.* **54**, 221
 Smak, J.: 2007, *Acta Astron.* **57**, 87
 Smak, J.: 2008 a, *Acta Astron.* **58**, 55
 Smak, J.: 2008 b, *Acta Astron.* **58**, 65