

A Search for Photometrically Variable Magnetic CP Stars in Sky Survey Data

S. Hümmerich^{1,2}, K. Bernhard^{1,2} and E. Paunzen³

¹ *American Association of Variable Star Observers (AAVSO),
Cambridge, MA, 02138, USA (E-mail: ernham@rz-online.de)*

² *Bundesdeutsche Arbeitsgemeinschaft für Veränderliche Sterne e.V. (BAV),
D-12169 Berlin, Germany*

³ *Department of Theoretical Physics and Astrophysics, Masaryk University,
CZ-611 37 Brno, Czech Republic*

Received: November 10, 2017; Accepted: November 12, 2017

Abstract. The magnetic chemically peculiar (mCP) stars of the upper main sequence are natural laboratories well-suited for the investigation of such diverse phenomena as e.g. atmospheric structure, magnetic fields and stellar rotation. We here present our efforts towards increasing the sample of known mCP stars by using publicly available photometric time series data, which has already led to the discovery of more than 850 photometrically variable mCP stars and candidates.

Key words: stars: chemically peculiar – variables: general – surveys

1. Introduction

Chemically peculiar (CP) stars make up about 10% of upper main-sequence stars (spectral types early B to early F). They are characterized by peculiar atmospheric abundances that differ significantly from the solar pattern. The CP2 (Bp/Ap) stars and the CP4 (helium-weak) stars (also referred to collectively as the magnetic CP or mCP stars) are set apart by the presence of strong, stable and globally-organized magnetic fields (Babcock, 1947; Aurière et al., 2007). They exhibit a non-uniform distribution of chemical elements on their surfaces, which leads to spectral and photometric variability with the rotation period. mCP stars exhibiting photometric variability are traditionally referred to as α^2 Canum Venaticorum (ACV) variables.

During recent decades, automated sky surveys have advanced to become the dominant data source in astronomy. An increasing number of surveys routinely produce high-quality photometric time-series data well suited to the investigation of stellar variability. We have taken advantage of these data in the identification and investigation of new ACV variables. Two different approaches were taken: (i) mCP stars from the Catalogue of Ap, HgMn, and Am stars (Renson & Manfroid, 2009) were selected and cross-matched with archival sky survey data, and (ii) a search for the typical variability pattern of ACV variables was carried

out among the light curves of early-type (spectral types B/A or colour-selected) variable stars of undetermined type in the AAVSO International Variable Star Index (VSX Watson et al., 2006).

2. Results

We here present an overview over the present state of our project. Following the methodology outlined above, we have established light variability in a large number of mCP stars not hitherto known as photometric variables (Table 1).

Table 1. New photometrically variable ACV stars and candidates.

Data Source	#ACVs	Ref.
All Sky Automated Survey (ASAS)	316	Bernhard et al. (2015a)
	357	Hümmerich et al. (2016)
SuperWASP (SWASP)	80	Bernhard et al. (2015b)
KELT Transit Survey	49	[unpublished]
Kepler (also K2 mission)	>50	[unpublished]
total of new variables	852+	

Among our sample, there are 14 eclipsing binary systems hosting an mCP star component. A lack of close binaries is generally observed among mCP stars; therefore, this finding is of great interest. Furthermore, we have identified several stars exhibiting variability on time-scales typical of δ Scuti variables (periods between about 15 minutes and 5 hours). While most of these stars are actually non-magnetic CP1 (Am) stars, several mCP star candidates are present. Spectroscopic verification, however, is needed and under way. In addition, some objects of special interest have been found, such as HD 66051, which has been identified as an eclipsing binary system hosting a highly-peculiar CP3(HgMn)-related star. Obvious out-of-eclipse variability is present in the light curve and interpreted as a signature of surface inhomogeneities (Niemczura et al., 2017).

Some future research fields deriving from our main project are outlined in the following: (i) extension of the search for new ACV variables to other photometric survey databases, (ii) spectroscopic confirmation of the ‘photometric’ ACV candidates (first results have shown that this might be a viable and efficient means of discovering new mCP stars), (iii) datamining of spectroscopic survey archives (LAMOST and SDSS) for new mCP stars, (iv) search for very long-period ($P > 1$ yr) ACV variables in survey data, and (v) investigation into the properties of the mCP stars that do not show photometric variability in the accuracy limit of the employed sky survey data.

References

- Aurière, M., Wade, G. A., Silvester, J., et al. 2007, *Astron. Astrophys.*, **475**, 1053
- Babcock, H. W. 1947, *Astrophys. J.*, **105**, 105
- Bernhard, K., Hümmerich, S., Otero, S., & Paunzen, E. 2015a, *Astron. Astrophys.*, **581**, A138
- Bernhard, K., Hümmerich, S., & Paunzen, E. 2015b, *Astron. Nachr.*, **336**, 981
- Hümmerich, S., Paunzen, E., & Bernhard, K. 2016, *Astron. J.*, **152**, 104
- Niemczura, E., Hümmerich, S., Castelli, F., et al. 2017, *Sci. Rep.*, **7**, 5906
- Renson, P. & Manfroid, J. 2009, *Astron. Astrophys.*, **498**, 961
- Watson, C. L., Henden, A. A., & Price, A. 2006, *Society for Astronomical Sciences Annual Symp.*, **25**, 47



Shinsuke Takasao, Christian Fendt and Wouter Vlemmings