



A LETTERS JOURNAL EXPLORING
THE FRONTIERS OF PHYSICS

OFFPRINT

Mermin's pentagram as an ovoid of $PG(3, 2)$

M. SANIGA and P. LÉVAY

EPL, **97** (2012) 50006

Please visit the new website
www.epljournal.org



A LETTERS JOURNAL EXPLORING
THE FRONTIERS OF PHYSICS

AN INVITATION TO SUBMIT YOUR WORK

www.epljournal.org

The Editorial Board invites you to submit your letters to EPL

EPL is a leading international journal publishing original, high-quality Letters in all areas of physics, ranging from condensed matter topics and interdisciplinary research to astrophysics, geophysics, plasma and fusion sciences, including those with application potential.

The high profile of the journal combined with the excellent scientific quality of the articles continue to ensure EPL is an essential resource for its worldwide audience. EPL offers authors global visibility and a great opportunity to share their work with others across the whole of the physics community.

Run by active scientists, for scientists

EPL is reviewed by scientists for scientists, to serve and support the international scientific community. The Editorial Board is a team of active research scientists with an expert understanding of the needs of both authors and researchers.



IMPACT FACTOR
2.753*
* As ranked by ISI 2010

www.epljournal.org

IMPACT FACTOR

2.753*

* As listed in the ISI® 2010 Science Citation Index Journal Citation Reports

OVER

500 000

full text downloads in 2010

30 DAYS

average receipt to online publication in 2010

16 961

citations in 2010
37% increase from 2007

“We’ve had a very positive experience with EPL, and not only on this occasion. The fact that one can identify an appropriate editor, and the editor is an active scientist in the field, makes a huge difference.”

Dr. Ivar Martin

Los Alamos National Laboratory,
USA

Six good reasons to publish with EPL

We want to work with you to help gain recognition for your high-quality work through worldwide visibility and high citations.

- 1 Quality** – The 40+ Co-Editors, who are experts in their fields, oversee the entire peer-review process, from selection of the referees to making all final acceptance decisions
- 2 Impact Factor** – The 2010 Impact Factor is 2.753; your work will be in the right place to be cited by your peers
- 3 Speed of processing** – We aim to provide you with a quick and efficient service; the median time from acceptance to online publication is 30 days
- 4 High visibility** – All articles are free to read for 30 days from online publication date
- 5 International reach** – Over 2,000 institutions have access to EPL, enabling your work to be read by your peers in 100 countries
- 6 Open Access** – Articles are offered open access for a one-off author payment

Details on preparing, submitting and tracking the progress of your manuscript from submission to acceptance are available on the EPL submission website www.epletters.net.

If you would like further information about our author service or EPL in general, please visit www.epljournal.org or e-mail us at info@epljournal.org.

EPL is published in partnership with:



European Physical Society



Società Italiana di Fisica



EDP Sciences

IOP Publishing



IOP Publishing




A LETTERS JOURNAL
EXPLORING THE FRONTIERS
OF PHYSICS

EPL Compilation Index


www.epljournal.org



Biaxial strain on lens-shaped quantum rings of different inner radii, adapted from **Zhang et al** 2008 *EPL* **83** 67004.



Artistic impression of electrostatic particle-particle interactions in dielectrophoresis, adapted from **N Aubry and P Singh** 2006 *EPL* **74** 623.



Artistic impression of velocity and normal stress profiles around a sphere that moves through a polymer solution, adapted from **R Tuinier, J K G Dhont and T-H Fan** 2006 *EPL* **75** 929.

Visit the EPL website to read the latest articles published in cutting-edge fields of research from across the whole of physics.

Each compilation is led by its own Co-Editor, who is a leading scientist in that field, and who is responsible for overseeing the review process, selecting referees and making publication decisions for every manuscript.

- Graphene
- Liquid Crystals
- High Transition Temperature Superconductors
- Quantum Information Processing & Communication
- Biological & Soft Matter Physics
- Atomic, Molecular & Optical Physics
- Bose-Einstein Condensates & Ultracold Gases
- Metamaterials, Nanostructures & Magnetic Materials
- Mathematical Methods
- Physics of Gases, Plasmas & Electric Fields
- High Energy Nuclear Physics

If you are working on research in any of these areas, the Co-Editors would be delighted to receive your submission. Articles should be submitted via the automated manuscript system at www.epletters.net

If you would like further information about our author service or EPL in general, please visit www.epljournal.org or e-mail us at info@epljournal.org



IOP Publishing

Image: Ornamental multiplication of space-time figures of temperature transformation rules (adapted from T. S. Bíró and P. Ván 2010 *EPL* **89** 30001; artistic impression by Frédérique Swist).

Mermin's pentagram as an ovoid of $PG(3, 2)$

M. SANIGA^{1(a)} and P. LÉVAY²

¹ *Astronomical Institute, Slovak Academy of Sciences - SK-05960 Tatranská Lomnica, Slovak Republic, EU*

² *Department of Theoretical Physics, Institute of Physics, Budapest University of Technology and Economics H-1521 Budapest, Hungary, EU*

received 2 December 2011; accepted 26 January 2012
published online 2 March 2012

PACS 02.40.Dr – Euclidean and projective geometries

PACS 03.65.Aa – Quantum systems with finite Hilbert space

PACS 03.67.-a – Quantum information

Abstract – Mermin's pentagram, a specific set of ten three-qubit observables arranged in quadruples of pairwise commuting ones into five edges of a pentagram and used to provide a very simple proof of the Kochen-Specker theorem, is shown to be isomorphic to an ovoid (elliptic quadric) of the three-dimensional projective space of order two, $PG(3, 2)$. This demonstration employs properties of the real three-qubit Pauli group embodied in the geometry of the symplectic polar space $W(5, 2)$ and rests on the facts that: 1) the four observables/operators on any of the five edges of the pentagram can be viewed as points of an affine plane of order two, 2) all the ten observables lie on a hyperbolic quadric of the five-dimensional projective space of order two, $PG(5, 2)$, and 3) that the points of this quadric are in a well-known bijective correspondence with the lines of $PG(3, 2)$.

Copyright © EPLA, 2012

Introduction. – In 1993, Mermin [1] proposed two ingenious and remarkably simple geometrical proofs of the Kochen-Specker theorem. The first one, which became known as Mermin's (magic) square, employs a set of nine elements/observables of the generalized Pauli group of *two*-qubits. These nine observables are placed at the vertices of a 3×3 grid and form six sets of three pairwise commuting elements, lying along three horizontal and three vertical lines, each observable thus pertaining to two such sets. The observables are selected in such a way that the product of their triples in five of the six sets is $+I$, whilst in the remaining set it is $-I$, I being the identity matrix. The second proof, known as Mermin's (magic) pentagram, uses a set of ten elements of the *three*-qubit Pauli group. Here, the ten observables form five sets of four mutually commuting elements placed along the five edges of a pentagram. Again, each observable belongs to two such sets ("contexts") and the product of fours in any given set is $+I$ except for one where it yields $-I$.

Soon after the properties of N -qubit Pauli groups were found to be fully encoded in the geometry of the symplectic polar spaces of rank N and order two, $W(2N - 1, 2)$ [2–5], the Mermin (magic) square could readily be ascribed to a neat finite geometrical meaning, namely as: a special kind of geometric hyperplane of $W(3, 2)$ [2,3],

a hyperbolic quadric in $PG(3, 2)$ [4], or a projective line over the direct product of two smallest Galois fields, $P_1(GF(2) \times GF(2))$ [6]. These last findings repeatedly turned out to be of great physical importance, for example, in revealing a fascinating finite-geometrical meaning of the $E_{6(6)}$ -symmetric black hole entropy formula of $D = 5$ supergravity theories [7]. In the present paper, employing basic properties of the finite symplectic polar space behind the three-qubit Pauli group, we shall show that the structure of the Mermin pentagrammatic configuration can similarly be recast in terms of another well-known object of finite geometry.

Mermin's pentagram, affine plane of order two, the Klein quadric, the Klein correspondence and an ovoid of $PG(3, 2)$. – Our starting point is a particular copy of the Mermin pentagram depicted in fig. 1, left; here I is the 2×2 identity matrix, $X \equiv \sigma_x$, $Z \equiv \sigma_z$, and, e.g., ZZZ is a shorthand for $Z \otimes Z \otimes Z$. The set of ten three-qubit operators we adopted is, however, not that of Mermin [1], but that of Aravind [8], who was motivated by the paper by Kernaghan and Peres [9]. The reason is that all the ten matrices are real and can thus be viewed as a subset of the *real* three-qubit Pauli group.

In order to "decipher" the structure of our Mermin's pentagram one has to invoke certain aspects of the finite geometric interpretation of the real three qubit Pauli

^(a)E-mail: msaniga@astro.sk




Fig. 1: (Colour on-line) Left: an illustration of the Mermin pentagram. The four three-qubit observables along any edge are pairwise commuting; the product of those along the horizontal (red) edge is $-III$, while those along any other edge multiply to $+III$. Right: a picture of the finite geometric configuration behind the Mermin pentagram: the five edges of the pentagram correspond to five copies of the affine plane of order two, sharing pairwise a single point.




Fig. 2: An illustration of the case (left) where no three of the four observables (represented by empty circles) pertaining to an edge are collinear in the associated Fano plane; they form an affine plane of order two (right).

group [2–4]. Omitting the identity and disregarding signs, 63 elements of this group correspond to 63 points of the symplectic polar space $W(5, 2)$, and maximal subsets of pairwise commuting elements of the group have their counterparts in maximal totally isotropic subspaces of $W(5, 2)$, which are Fano planes. As each Fano plane has seven points, any maximal subset of mutually commuting operators is of cardinality seven. Each edge from our Mermin’s pentagram, obviously, selects from its ambient Fano plane only four points. There are two possibilities for such a selection; either no three of these four points are collinear (fig. 2), or three of them lie on a line in the corresponding Fano plane (fig. 3). The second possibility can, however, be readily disregarded as the four observables on any edge have obviously the same footing. This implies that each edge of the pentagram is a copy of the affine plane of order two and the pentagram framework can be reformulated as five affine planes of order two sharing pairwise a single point, no three being on the same point —see fig. 1, right. Next we employ the fact that we are dealing with the real three-qubit Pauli group. In such a case [4] the structure of the symplectic polar space $W(5, 2)$ is refined in terms of the orthogonal polar space $Q^+(5, 2)$ (which is nothing but the famous Klein quadric), the points of which correspond to the *symmetric*




Fig. 3: An illustration of a hypothetical case where three of the four observables would lie on a line of the Fano plane.




Fig. 4: (Colour on-line) The $PG(3, 2)$ -image of the Mermin pentagram under the Klein correspondence. The edges of the pentagram correspond to the points of an ovoid/elliptic quadric, and its ten observables map into the lines joining the points in pairs.

operators/elements of the group. As all ten observables of our pentagram are symmetric, they must lie on $Q^+(5, 2)$. Now, the generators of $Q^+(5, 2)$ (*i.e.*, maximal subspaces fully lying on $Q^+(5, 2)$) are Fano planes and there are two systems of them, each having 15 members. Any two distinct planes from the same system share a point, whilst two planes from different systems are either disjoint, or have a line in common (see, *e.g.*, [10,11]). It then follows that our five affine planes must all originate from Fano planes of the same system.

As a final step, we employ the famous Klein correspondence between the points of the Klein quadric $Q^+(5, 2)$ and the lines of $PG(3, 2)$ (see, *e.g.*, table 15.10 of [10] for more details). Under this correspondence, the seven points in a plane of one system of $Q^+(5, 2)$ correspond to the seven lines through a point of $PG(3, 2)$, and those of a plane of the other system to the seven lines lying in a plane of $PG(3, 2)$; an affine plane of order two lying on $Q^+(5, 2)$ will then have for its $PG(3, 2)$ -image either four lines through a point, no three coplanar, or four lines in a plane, no three concurrent. Adopting the former view, we thus find that the $PG(5, 2)$ -configuration depicted in fig. 1, right, has for its $PG(3, 2)$ -counterpart the set of five points, no three collinear and no four lying in the same plane, that is, a copy of an *ovoid* (or, elliptic quadric) (see, *e.g.*, [10]). The ten three-qubit Pauli matrices of the pentagram are thus represented by ten lines joining the five points of the ovoid in pairs —as illustrated in fig. 4.

We started this section with the observation that the four observables sharing an edge of the pentagram do not represent a maximal commuting set in the corresponding three-qubit Pauli group. We also stressed that any such maximal set consists of seven elements carried by a Fano plane of $W(5, 2)$. It is, therefore, instructive to see explicitly the projective closures of the edges (up to signs):

$$\begin{aligned} \{ZZI, ZIZ, IZZ\} & \quad (\text{for green edge}), \\ \{XXI, XIZ, IXZ\} & \quad (\text{for blue edge}), \\ \{XIX, IZX, XZI\} & \quad (\text{for violet edge}), \\ \{IXX, ZXI, ZIX\} & \quad (\text{for yellow edge}), \\ \{IYY, YIY, YYI\} & \quad (\text{for red edge}), \end{aligned}$$

where $Y \equiv i\sigma_y$. One can readily check that, up to a sign, the product of each of the five triples of observables is the identity matrix and so each of them represents indeed a line in the associated Fano plane (which is highlighted in fig. 2, left, by three big bullets). One further notes that only the observables of the red-edge projective closure contain the matrix Y , which reflects the fact that this edge stands on a different footing than the other four.

Conclusion. – We have shown that the Mermin pentagrammatic framework of ten three-qubit observables, which furnishes a very economic and elegant proof of the Kochen-Specker theorem, can be viewed as a distinguished object of finite geometry, namely an ovoid (elliptic quadric) of PG(3, 2); the five edges of the pentagram correspond to the five points of the ovoid and its ten vertices/observables are represented by the lines joining pairs of points of the ovoid. This finding may serve as another justification of our firm belief that all important sets of operators/observables associated with finite-dimensional Hilbert spaces are underlaid by notable objects of finite geometry. In the light of this paper, a particularly interesting and challenging task

would be to look for higher-rank analogues of Mermin's configuration(s). Already the next case in the hierarchy, $N = 4$, deserves serious attention. This is mainly because an associated hyperbolic quadric $Q^+(7, 2)$, the locus of symmetric elements of the four-qubit Pauli group, is unusual in that it admits a graph automorphism of order three known as a triality that swaps its points and two systems of generators, and preserves the set of totally singular lines (see, e.g., [12]).

This work was partially supported by the VEGA grant agency projects 2/0092/09 and 2/0098/10. We are extremely grateful to our friend P. PRACNA for electronic versions of the figures.

REFERENCES

- [1] MERMIN N. D., *Rev. Mod. Phys.*, **65** (1993) 803.
- [2] SANIGA M. and PLANAT M., *Adv. Stud. Theor. Phys.*, **1** (2007) 1.
- [3] PLANAT M. and SANIGA M., *Quantum Inf. Comput.*, **8** (2008) 127.
- [4] HAVLICEK H., ODEHNAL B. and SANIGA M., *SIGMA*, **5** (2009) 096.
- [5] THAS K., *EPL*, **86** (2009) 60005.
- [6] SANIGA M., PLANAT M. and PRACNA P., *Theor. Math. Phys.*, **155** (2008) 905.
- [7] LÉVAY P., SANIGA M., VRANA P. and PRACNA P., *Phys. Rev. D*, **79** (2009) Art. No. 084036.
- [8] ARAVIND P. K., *Mermin's Pentagram and Bell's Theorem* (arXiv:quant-ph/0104138).
- [9] KERNAGHAN M. and PERES A., *Phys. Lett. A*, **198** (1995) 1.
- [10] HIRSCHFELD J. W. P., *Finite Projective Spaces of Three Dimensions* (Clarendon, Oxford) 1985.
- [11] CONWELL G. M., *Ann. Math.*, **11** (1910) 60.
- [12] TITS J., *Inst. Hautes Etudes Sci. Publ. Math.*, **2** (1959) 13.